

Quark Number Susceptibility, Thermodynamic Sum Rule, and Hard Thermal Loop Approximation

Purnendu Chakraborty¹, Munshi G. Mustafa^{1,2}, and Markus H. Thoma³

¹*Theory Group, Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700 064, India*

²*Institut für Theoretische Physik, Universität Giessen, 35392 Giessen, Germany*

³*Centre for Interdisciplinary Plasma Science, Max-Planck-Institut für extraterrestrische Physik, P.O. Box 1312, 85741 Garching, Germany*

The quark number susceptibility, associated with the conserved quark number density, is closely related to the baryon and charge fluctuations in the quark-gluon plasma, which might serve as signature for the quark-gluon plasma formation in ultrarelativistic heavy-ion collisions. Beside QCD lattice simulations, the quark number susceptibility has been calculated recently using a resummed perturbation theory (Hard Thermal Loop resummation). In the present work we show, based on general arguments, that the computation of this quantity neglecting Hard Thermal Loop vertices contradicts the Ward identity and violates the thermodynamic sum rule following from the quark number conservation. We further show that the Hard Thermal Loop perturbation theory is consistent with the thermodynamic sum rule.

Dynamical properties of a many particle system can be investigated by employing an external probe, which disturbs the system only slightly in its equilibrium state, and by measuring the response of the system to this external perturbation. A large number of experiments belong to this category [1] such as studies of various line-shapes, acoustic attenuation, and transport behavior. In all these experiments, one probes the dynamical behavior of the spontaneous fluctuations in the equilibrium state. In general, spontaneous fluctuations are related to correlation functions. They reflect the symmetries of the system, which provide important inputs for quantitative calculations of complicated many-body systems [2].

Recently, screening and fluctuations of conserved quantities, such as baryon number and electric charge, have been proposed as a probe of the quark-gluon plasma (QGP) formation in ultrarelativistic heavy-ion collisions [3–10]. These fluctuations are closely related to the quark number susceptibility (QNS) [3,7], which measures the response of the number density to an infinitesimal change of the quark chemical potential.

The QNS has been considered within lattice gauge theory [11–13], in perturbative QCD [14], as well as effective models of QCD [3,7–10]. Recently it has also been computed within the 2-loop approximately self-consistent Φ -derivable Hard Thermal Loop (HTL) resummation in Ref. [15] and within one-loop HTL perturbation theory (HTLpt) in Ref. [16]. Whereas the result of Ref. [16] is in good agreement with lattice results, Ref. [15] overestimates the lattice data. Note, however, that the entropy and pressure obtained within the 2-loop approximately

self-consistent Φ -derivable method agree well with lattice data [17].

In Ref. [18] possible explanations for the different results of the various HTL approaches have been discussed. As we will argue here, the neglect of the HTL vertices violates the Ward identities and the thermodynamic sum rule following from the conservation of the quark number [1,3,7]. We show, based on general arguments, that the HTLpt, on the other hand, is consistent with the thermodynamic sum rule.

The HTL resummation technique allows a systematic gauge invariant treatment of gauge theories at finite temperature and chemical potential taking into account medium effects such as Debye screening, effective quark masses, and Landau damping [19]. The generating functional which generates the HTL Green functions between a quark pair and any number of gauge bosons can be written [19,20] as

$$\delta\mathcal{L} = m_q^2 \bar{\psi} \left\langle \frac{K}{K \cdot D} \right\rangle \psi, \quad (1)$$

where K^μ is a light like four-vector, D_μ is the covariant derivative, m_q is thermal quark mass, and $\langle \rangle$ is the average over all possible directions over loop momenta. This functional is gauge symmetric and nonlocal and leads to the following Dirac equation

$$\not{D}\psi = \Sigma\psi + \Gamma_\mu A^\mu\psi + A^\mu\Gamma_{\mu\nu}A^\nu\psi + \dots, \quad (2)$$

where we have suppressed the color index. In the HTL approximation the 2-point function, $\Sigma \sim gT$, (quark self-energy) is of the same order as the tree level one, $S_0^{-1}(K) \sim K \sim gT$ (in the weak coupling limit $g \ll 1$), if the external momenta are soft, *i.e.* of the order gT . The 3-point function, *i.e.*, the effective quark-gauge boson vertex, is given by $g\Gamma_\mu = g(\gamma_\mu + \delta\Gamma_\mu)$, where $\delta\Gamma_\mu$ is the HTL correction. The 4-point function, $g^2\Gamma_{\mu\nu}$, does not exist at the tree level and only appears within the HTL approximation [19]. These N -point functions, which are complicated functions of momenta and energies, are interrelated by Ward identities. This has important consequences for the QNS (and other quantities) if one aims at a consistent perturbative expansion.

The HTLpt [21] is an extension of the screened perturbation theory [22] in such a way that it amounts to a reorganization of the usual perturbation theory by adding

and subtracting the nonlocal HTL term (1) in the action [19,23]. The added term together with the original QCD action is treated nonperturbatively as zeroth order whereas the subtracted one as perturbation. Equivalently, for calculating physical quantities this amounts to replace the bare propagators and vertices by resummed HTL Green functions [19]. In this way, interesting quantities of the QGP have been computed [21,24–31] within the HTLpt. However, HTLpt suffers from the fact that one uses HTL resummed Green functions also for hard momenta of the order T . A systematic description of physical quantities requires an explicit separation of hard ($\sim T$) and soft ($\sim gT$) scales [19]. Following this approach, a number of relevant physical quantities, such as signatures of the QGP, has consistently been calculated [32]. A convincing reason for studying the QNS in HTL approximation is the fact that this approach has also been used for the free energy of the QGP, where different implementations of the HTL method and their validity have been discussed. The QNS may serve as another test for these approaches.

The usual definition of the QNS for a given flavor which measures the response of the quark number density ρ_q to an infinitesimal change in the quark chemical potential $\mu_q + \delta\mu_q$ is given by [1,3,7]:

$$\chi_q(T) = \left. \frac{\partial \rho_q}{\partial \mu_q} \right|_{\mu_q=0}. \quad (3)$$

Since the quark (baryon) number is a conserved quantity, the QNS obeys a Kubo relation [1]

$$\chi_q(T) = \beta \int d^3x \langle j_0(0, \vec{x}) j_0(0, \vec{0}) \rangle, \quad (4)$$

where j_0 is the time component of the quark current $j_\mu = \bar{\psi} \gamma_\mu \psi$. The equivalence of the static susceptibility (3) and the static limit of the dynamical susceptibility (4) follows from [1]

$$\rho_q = \frac{1}{V} \frac{\text{Tr} [\mathcal{N}_q e^{-\beta(\mathcal{H} - \mu_q \mathcal{N}_q)}]}{\text{Tr} [e^{-\beta(\mathcal{H} - \mu_q \mathcal{N}_q)}]} = \frac{\langle \mathcal{N}_q \rangle}{V}, \quad (5)$$

with the quark number operator $\mathcal{N}_q = \int d^3x j_0(0, \vec{x})$ and volume V . The derivative with respect to μ_q in (5) according to (3) agrees with (4) *only if the number operator is a conserved quantity and hence commutes with the Hamiltonian $\mathcal{H}$* [1].

Introducing the vector meson correlator $S_{\mu\nu}(t, \vec{x}) = \langle j_\mu(t, \vec{x}) j_\nu(0, \vec{0}) \rangle$ and taking the Fourier transform of $S_{00}(0, \vec{x})$, the QNS according to (4) can be written as [33]

$$\begin{aligned} \chi_q(T) &= \beta \int d^3x \int \frac{d\omega}{2\pi} \int \frac{d^3p}{(2\pi)^3} e^{-i\vec{p} \cdot \vec{x}} S_{00}(\omega, p) \\ &= \beta \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi} S_{00}(\omega, 0). \end{aligned} \quad (6)$$

This equation is known as a *thermodynamic sum rule* as it relates the thermodynamic derivatives to the correlation

function due to the symmetry of the system. Examples of such sum rules for various systems can be found in Ref. [1]. The fluctuation - dissipation theorem relates S_{00} to the vector meson self-energy Π_{00} via [1]

$$S_{00}(\omega, p) = \frac{2}{\exp(-\beta\omega) - 1} \text{Im} \Pi_{00}(\omega, p). \quad (7)$$

An example for this self-energy within the 1-loop order with effective Green functions is shown in Fig.1, where the blobs indicate effective quark propagators and quark-meson vertices.

The time independence of $\mathcal{N}_q$, corresponding to the continuity equation $\partial_\mu j^\mu = 0$, implies that the quantity $\int d^3x S_{00}(0, \vec{x})$ in (4) is time independent. This is guaranteed if $S_{00}(\omega, 0) \propto \delta(\omega)$ [1].




Fig.1: The 1-loop vector meson self-energy diagrams.

In usual perturbation theory the equivalence of (3) and (4) follows from the fact that the loop expansion agrees with an expansion in the coupling constant. For example, to lowest order the number density follows from the tadpole diagram (containing only a bare propagator and vertex) at finite temperature and chemical potential, which corresponds to the 1-loop polarization tensor of the vector correlator (7). For the HTL resummation the loop expansion and the coupling expansion are not equivalent as the first one includes higher order effects through resummation. This mixing can cause an incompleteness in physical quantities computed in a given loop order, as we shall discuss below.

Now we would like to discuss how the QNS can be calculated consistently within the HTL approximation. Our starting point is the most general expression of the quark self-energy [34], which enters the full quark propagator. In the rest frame of a medium in the chiral limit it reads

$$\Sigma(K) = -a(k_0, k) \not{K} - b(k_0, k) \gamma^0, \quad (8)$$

where the scalar quantities a and b are functions of the energy k_0 and of the magnitude of the spatial momentum, k , and $K = (k_0, \vec{k})$. The full propagator is given by

$$S_F^{-1}(K) = \not{K} - \Sigma(K) = (1 + a) \not{K} + b \gamma^0. \quad (9)$$

Using the helicity representation [29] it can also be written as

$$\begin{aligned} S_F(k_0, k) &= \frac{\gamma^0 - \hat{k} \cdot \vec{\gamma}}{2D_+(k_0, k)} + \frac{\gamma^0 + \hat{k} \cdot \vec{\gamma}}{2D_-(k_0, k)}, \\ D_\pm(k_0, k) &= (-k_0 \pm k)(1 + a) - b. \end{aligned} \quad (10)$$

Charge invariance demands $D_{\pm}(-k_0, k) = -D_{\mp}(k_0, k)$, which implies $a(-k_0, k) = a(k_0, k)$ and $b(-k_0, k) = -b(k_0, k)$. The zeros of $D_{\pm}$ describe the in-medium propagation or *quasiparticle* (QP) dispersion relation of a particle excitation with energy $\omega_{\pm}(k)$ and of a mode called plasmino with energy, $\omega_{-}(k)$, which exists only in the medium [29,35]. In addition, $D_{\pm}$ contains a discontinuous part corresponding to *Landau damping* (LD) if the quark self-energy has an imaginary part.

Using these general properties of the quark propagator, the spectral representation of the in-medium propagator (10) can be written as

$$\rho_{\pm}(\omega, k) = -\frac{1}{\pi} \text{Im} \frac{1}{D_{\pm}} = \mathcal{R}_{\pm}(\omega, k) \delta(\omega - \omega_{\pm}) + \mathcal{R}_{\mp}(-\omega, k) \delta(\omega + \omega_{\mp}) + \rho_{\pm}^{\text{dis}}(\omega, k) \quad , \quad (11)$$

where $\mathcal{R}_{\pm} = (dD_{\pm}/d\omega)^{-1}$ are the residues corresponding to the poles of the propagators, and $\rho_{\pm}^{\text{dis}}$ are discontinuities. In the HTL-approximation the discontinuities, given explicitly e.g. in Ref. [16], are located below the light cone ($k_0^2 < k^2$), where Landau damping occurs. We also note that for each pole the spectral function will involve a δ -function. The Ward identity relates the quark propagator to the quark-meson vertex via

$$(K_2 - K_1)_{\mu} \Gamma^{\mu}(-K_2, K_1; K_2 - K_1) = S_F^{-1}(K_2) - S_F^{-1}(K_1). \quad (12)$$

In the HTL approximation the effective vertices are discussed in Ref. [16].

Using the effective propagators and vertices given in (10) and (12), the contribution of the first diagram in Fig. 1 can be written as

$$\begin{aligned} \text{Im} \Pi_{00}^s(\omega, 0) &= 4N_f N_c \pi (1 - e^{\beta\omega}) \int \frac{d^3k}{(2\pi)^3} \int dx \int dy \\ &\times \delta(\omega - x - y) n_F(x) n_F(y) \left\{ (1 - f(x, y))^2 \right. \\ &\times \rho_+(x, k) \rho_-(y, k) + \frac{1}{\omega^2 \pi} [\text{Im} F_+(x, k) \\ &\times \rho_-(y, k) + \text{Im} F_-(y, k) \rho_+(x, k)] \left. \right\}, \quad (13) \end{aligned}$$

where $F_{\pm}(z, k) = D_{\pm}(z, k) + z \mp k$, z is the energy of the internal quarks, $f(x, y) = (x + y)/\omega$, N_f is the number of quark flavors, and N_c is the number of colors. The contribution of the tadpole diagram, $\text{Im} \Pi_{00}^t(\omega, 0)$, in Fig. 1 is identical apart from an opposite sign of the second term inside the curly braces in (13). Using the explicit form of the spectral functions (11), the total contribution of Fig. 1 is

$$\begin{aligned} \text{Im} \Pi_{00}(\omega, 0) &= [\text{Im} \Pi_{00}^s(\omega) + \text{Im} \Pi_{00}^t(\omega)] = 4N_f N_c \pi \\ &(1 - e^{\beta\omega}) \delta(\omega) \int \frac{d^3k}{(2\pi)^3} [\mathcal{R}_+^2(\omega_+, k) n_F(\omega_+) n_F(-\omega_+) \\ &+ \mathcal{R}_-^2(\omega_-, k) n_F(\omega_-) n_F(-\omega_-)] \quad , \quad (14) \end{aligned}$$

where the LD contributions explicitly vanish. $\text{Im} \Pi_{00}$ in (14) contains only the QP contributions following from the dispersion relations, $D_{\pm} = 0$, and the requirement for the *conservation law*, $S_{00} \propto \delta(\omega)$, is satisfied. This necessitates the use of effective 3- and 4-point functions and also the modification of the charge operator, as $\bar{\psi}(\gamma_0 + \delta\Gamma_0)\psi$, by inclusion of the effective 3-point function, in contrast to Ref. [15,18]. This non-local modification makes the charge operator different from that of (3) and (4) but is essential to satisfy the Ward identity which is related to the gauge invariance and conservation law. This implies that the number density has to be calculated with propagators and vertices related by Ward identities at finite T and μ_q .




Fig.2: The QNS in different approximations (see text) divided by the lowest order result (free QNS) as a function of T/T_c with $T_c/\Lambda_{\overline{MS}} = 0.49$ [36]. In each band the lower curve corresponds to the choice of the renormalization scale $\bar{\mu} = 2\pi T$ and the upper one to $\bar{\mu} = 4\pi T$. The squares represent recent lattice data [13].

Now the 1-loop QNS follows from combining (14) and (6)

$$\begin{aligned} \chi_q(T) &= 4N_f N_c \beta \int \frac{d^3k}{(2\pi)^3} [\mathcal{R}_+^2(\omega_+, k) n_F(\omega_+) n_F(-\omega_+) \\ &+ \mathcal{R}_-^2(\omega_-, k) n_F(\omega_-) n_F(-\omega_-)] \quad , \quad (15) \end{aligned}$$

which is identical with the one-loop HTLpt result [16], if the residues are obtained from the HTL resummed propagators. In this case the functions $\mathcal{R}_{\pm}$ in (15) are given by

$$\mathcal{R}_{\pm}(\omega_{\pm}, k) = \frac{\omega_{\pm}^2(k) - k^2}{2m_q^2} \quad (16)$$

with the HTL quark dispersion relations $\omega_{\pm}(k)$ and the HTL effective quark mass $m_q = g(T)T/\sqrt{6}$. The result (band between the solid line denoted by QP in

Fig. 2) agrees well with recent lattice data [13]. (Here we used the 2-loop result for the running coupling constant $g(T)$ in contrast to Ref. [16], where the 1-loop expression was adopted. For the renormalization scale $\bar{\mu}$ entering the running coupling constant, we choose two different values leading to the bands in Fig. 2.) However, it overcounts the leading order perturbative result (dotted band denoted by PLO in Fig. 2) given as $\chi_q/\chi_f = (1 - 2\alpha_s/\pi + \dots)$ [18]. We note that this can be cured by going to the calculation at 2-loop order. Recently, it has been shown that the 2-loop order calculation [27,28] for the thermodynamic potential reveals the correct inclusion of the leading order effects but happens to be very close to the 1-loop result.

The physical relevance of the plasmino branch has been discussed in connection with the dilepton production in relativistic heavy-ion collisions [29,35]. Using the HTL approach its contribution to the QNS, i.e. the second term of (15), amounts only to less than one percent of the total susceptibility. This is caused by the fact that within the HTL approximation the spectral strength of the plasmino branch vanishes exponentially for hard momenta, $k \gg gT$, and that we integrate in (15) over all momenta.

Now, employing only resummed propagators and bare vertices as in Ref. [15], the QNS is obtained from (13) by setting the vertex correction functions, $f(x, y)$ and $F_{\pm}(x, y)$, equal to zero. Then the QNS contains LD contributions coming from the discontinuity of the 2-point HTL function in addition to the QP contribution in (15). The result is shown in Fig. 2 by the dashed band denoted by QP+LD which clearly overestimates the lattice data. Although the charge operator is local in this approach [18], $S_{00}(\omega, 0)$ is not proportional to $\delta(\omega)$, implying that $d\rho_q/dt \neq 0$. This amounts to a violation of the Ward identity, i.e. gauge invariance, for the effective HTL Green functions and also sacrifices the conservation law and hence the thermodynamic sum rule. Note, however, that it has been argued that the gauge dependence of the 2-loop effective action within the Φ -derivable approximation affects only higher order contributions [37].

In summary, we have discussed that the QNS can be computed consistently within the HTLpt in order to satisfy the requirement of the thermodynamic sum rule. This requires to consider 3-point and 4-point HTL functions, related to the HTL propagator by Ward identities. As discussed above, the QNS within the one-loop order overestimates the PLO, which, however, can be cured by extending the calculation to 2-loop orders in HTLpt. This would, indeed, be a challenging as well as nontrivial task due to its complexity. However, if it turns out to be a small correction, e.g., like in the case of the quark-gluon thermodynamic potential, the 1-loop HTLpt result presented here, agreeing well with lattice data, would be a useful approximation.

Acknowledgment: M.G.M. is thankful to C. Greiner, S. Juchem, S. Leupold and A. Peshier for various illumi-

nating discussions and to the AvH Foundation and GSI (Darmstadt) for financial support. M.H.T. was supported by DLR (BMBF) under grant no. 50WM0038.

- [1] D. Forster, *Hydrodynamic fluctuation, broken symmetry and correlation functions* (Benjamin/Cummings, Melno Park, CA, 1975).
- [2] H. B. Callen and T. A. Welton, Phys. Rev. **122** (1961) 34; R. Kubo, J. Phys. Soc. Japan **12** (1957) 570.
- [3] L. McLerran, Phys. Rev. D **36** (1987) 3291.
- [4] M. Roland et al., (NA49 Collaboration), Nucl. Phys. **A638** (1998) 91c; H. Appelshäuser et al., (NA49 Collaboration), Phys. Lett. B **459** (1999) 679; J. G. Reid, (NA49 Collaboration), Nucl. Phys. **A661** (1999) 407c.
- [5] M. Asakawa, U. Heinz, and B. Müller, Phys. Rev. Lett. **85** (2000) 2072.
- [6] S. Jeon and V. Koch, Phys. Rev. Lett. **85** (2000) 2076.
- [7] T. Hatsuda and T. Kunihiro, Phys. Rep. **247** (1994) 221; T. Kunihiro, Phys. Lett. B **271** (1991) 395.
- [8] G. E. Brown and M. Rho, Phys. Rept. **269** (1996) 333.
- [9] E. Shuryak, Rev. Mod. Phys. **65** (1996) 1.
- [10] M. Prakash and I. Zahed, Phys. Rev. Lett. **69** (1992) 3282; M. Prakash, R. Rapp, J. Wambach and I. Zahed, Phys. Rev. C **65** (2002) 034906.
- [11] S. Gottlieb et al., Phys. Rev. Lett. **59** (1987) 2247; Phys. Rev. D **38** (1988) 2888; R. Gavai, J. Potvin, and S. Sanielevici, Phys. Rev. D **40** (1989) R2743.
- [12] R. Gavai and S. Gupta, Phys. Rev. D **64** (2001) 074506.
- [13] R. V. Gavai and S. Gupta, Phys. Rev. D **65** (2002) 094515; R. V. Gavai, S. Gupta, and P. Majumdar, Phys. Rev. D **65** (2002) 054506.
- [14] A. Vuorinen, Phys. Rev. D **67** (2003) 074032.
- [15] J.-P. Blaizot, E. Iancu, and A. Rebhan, Phys. Lett. B **523** (2001) 143.
- [16] P. Chakroborty, M. G. Mustafa, and M. H. Thoma, Eur. Phys. C **23** (2002) 591.
- [17] J.-P. Blaizot, E. Iancu, and A. Rebhan, Phys. Rev. Lett. **83** (1999) 2906; Phys. Lett. B **470** (1999) 181; Phys. Rev. D **63** (2001) 065003.
- [18] J.-P. Blaizot, E. Iancu, and A. Rebhan, Eur. Phys. C **27** (2003) 433.
- [19] E. Braaten and R. D. Pisarski, Nucl. Phys. **B337** (1990) 569; Phys. Rev. Lett. **64** (1990) 1338.
- [20] E. Braaten and R. D. Pisarski, Phys. Rev. D **45** (1992) R1827.
- [21] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. Lett. **83** (1999) 2139.
- [22] F. Karsch, A. Patkós, and P. Petreczky, Phys. Lett. **B401** (1997).
- [23] J. Frenkel, and J. C. Taylor, Nucl. Phys. **B334** (1990) 199.
- [24] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. D **61** (2000) 014017.
- [25] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. D **61** (2000) 074016.
- [26] J. O. Andersen and M. Strickland, Phys. Rev. D **66** (2002) 105001.
- [27] J. O. Andersen, E. Braaten, E. Petitgirard and M. Strickland, Phys. Rev. D **66** (2002) 085016.
- [28] J. O. Andersen, E. Petitgirard and M. Strickland, (hep-

ph/0302069).

- [29] E. Braaten, R. D. Pisarski, and T. C. Yuan, Phys. Rev. Lett. **64** (1990) 2242.
- [30] P. Chakraborty, M. G. Mustafa, and M. H. Thoma, Phys. Rev. D **67** (2003) 114004.
- [31] F. Karsch, M. G. Mustafa, and M. H. Thoma, Phys. Lett. B **497** (2001) 249.
- [32] M.H. Thoma, in *Quark-Gluon Plasma 2*, ed.: R.C. Hwa (World Scientific, Singapore, 1995) p.51.
- [33] A. Das, *Finite Temperature Field Theory* (World Scientific, Singapore, 1997) p. 102.
- [34] H. A. Weldon, Phys. Rev. D **26** (1982) 2789.
- [35] A. Peshier and M. H. Thoma, Phys. Rev. Lett. **84** (2000) 841.
- [36] S. Gupta, Phys. Rev. D **64** (2001) 034507.
- [37] A. Arrizabalaga and J. Smit, Phys. Rev. D **66** (2002) 065014.